Question 1 1 out of 1 points Correct Before performing an abdominal examination, the examiner should:

Selected Answer: have the patient empty his or her bladder.

Question 2 1 out of 1 points Correct The most common form of birth trauma of the scalp is:

Selected Answer: caput succedaneum.

Question 3 1 out of 1 points Correct Unusual white areas on the skin may be due to:

Selected Answer: C. vitiligo.

Question 4 1 out of 1 points Correct Mrs. G. reports an increase in her alcohol intake over the past 5 years. To screen her for problem drinking, you would use the:

Selected Answer: CAGE questionnaire.

Question 5 1 out of 1 points Correct

Mr. Franklin is speaking with you, the health care provider, about his respiratory problem. Mr. Franklin says, "I've had this cough for 3 days, and it's getting worse." You reply, "Tell me more about your cough." Mr. Franklin states, "I wish I could tell you more. That's why I'm here. You tell me what's wrong!" Which caregiver response would be most appropriate for enhancing communication?

Selected Answer:

After 3 days, you're tired of coughing. Have you had a fever?

Question 6 1 out of 1 points Correct Expected hair distribution changes in older adults include:

Selected Answer: increased terminal hair follicles to the tragus of men's ears.

Question 7 1 out of 1 points Correct You have just completed a skin assessment on Mr. Baker. During your assessment, you have transilluminated a skin lesion. During the physical examination, you know that skin lesions are transilluminated to distinguish:

Selected Answer: C. fluid-filled from solid cysts or masses.

Question 8 1 out of 1 points Correct Percussing at the right midclavicular line, below the umbilicus, and continuing upward is the correct technique for locating the:

Selected Answer: lower liver border.

Question 9 1 out of 1 points Correct When you are questioning a patient regarding alcohol intake, she tells you that she is only a social drinker. Which initial response is appropriate?

Selected Answer: