

QUESTION 1

1. Select the neurotransmitter involved in obsessive-compulsive disorder (OCD) in children and adolescents based on the common comorbidity of tic disorders.

- A Norepinephrine
- B GABA
- C Dopamine
- D Acetylcholine

1 points

QUESTION 2

1. Select the class of drugs most effective in treating pain disorder.

- A Opioid analgesics
- B TCA and SSRI antidepressants
- C Beta-adrenergic blockers
- D Mood stabilizers

1 points

QUESTION 3

1. Select the most commonly occurring eating disorder.

- A Anorexia nervosa
- B Bulimia nervosa
- C Purging
- D Binge

1 points

QUESTION 4

1. Select the disorder in which inflicting injury to self or others is common.

- A Pain

- B Conversion
- C Somatic symptom
- D Factitious

1 points

QUESTION 5

1. Select **two** imaging methods used to study anxiety disorders.

- A CT
- B TMS
- C MRI
- D EKG

1 points

QUESTION 6

1. Select the **two** factors that do not cause dissociative amnesia.

- A Sexual abuse
- B Substance abuse
- C Surgical pain
- D Partner betrayal

1 points

QUESTION 7

1. Select the correct epidemiologic fact about major depression in children and adolescents.

- A There is a 20% incidence by age 18
- B The incidence is equal in adolescent females and males
- C There is a decreased incidence after 20
- D The incidence is five- to ten-fold greater in children with a parent or sibling with major depression